



Road expansion: A challenge to conservation of mammals, with particular emphasis on the endangered Asiatic cheetah in Iran

Alireza Mohammadi^a, Kamran Almasieh^b, Anthony P. Clevenger^c, Faezeh Fatemizadeh^a, Ali Rezaei^a, Houman Jowkar^d, Mohammad Kaboli^{a,*}

^a Department of Environmental Sciences, Faculty of Natural Resources, University of Tehran, Karaj, Iran

^b Department of Nature Engineering, Ramin Agriculture and Natural Resources University of Khuzestan, Mollasani, Iran

^c Western Transportation Institute, Montana State University, Bozeman, MT 59717 USA

^d Conservation of the Asiatic Cheetah Project (CACCP), Tehran, Iran

ARTICLE INFO

Keywords:

Asiatic cheetah
Distance method
Kernel density estimation
Wildlife vehicle collisions

ABSTRACT

Anthropogenic activities, including road expansion, are one of the main drivers of biodiversity loss in Iran. Central and northeastern Iran have been among the most vulnerable areas where expanding anthropogenic activities (in particular construction of road networks) have come into conflict with conservation and management of the endangered Asiatic cheetah (*Acinonyx jubatus venaticus*), along with other large mammals. The present study aimed to determine hotspot locations along an extremely high-risk road for mammals in northeast Iran (Touran Biosphere Reserve [TBR]) and propose mitigation measures for mammals such as the Asiatic cheetah. Using a spatially-explicit algorithm to estimate collision incidences, we adopted the kernel density estimation (KDE) and the distance method with respect to EDGE (evolutionarily distinct and globally endangered) and home range values for all locations. Also, a habitat suitability map was prepared to create habitat patches and applied to corridor modeling for the Asiatic cheetah. We investigated locations of 73 wildlife–vehicle collisions (WVCs) and crossing data from 2005–2016, that included Persian gazelles (*Gazella subgutturosa*), Asiatic cheetahs, striped hyenas (*Hyaena hyaena*), golden jackals (*Canis aureus*), red fox (*Vulpes vulpes*), European hare (*Lepus europaeus*), caracal (*Caracal caracal*), and grey wolves (*Canis lupus*). Our results showed that, based on the two methods, hotspot locations and the Asiatic cheetah corridor coincided. The corridor between TBR and Miandasht Wildlife Refuge was illustrated. This corridor is about 55 km length and 656 km² area, which connects two population patches within these protected areas. Asiatic cheetah vehicle collisions mostly occurred where a road crossed this corridor at the border of the TBR. The mitigation strategies proposed in this study for large mammals, particularly the Asiatic cheetah, are as follows: retrofit and installation of culverts in hotspots; installation of fences for crossing carnivores; and, roadside vegetation clearance in critical seasons for the Persian gazelles along the Semnan-Mashhad road.

1. Introduction

Human–wildlife conflicts including wildlife–vehicle collisions (WVCs) present significant challenges to conservation and management of mammals in human-dominated landscapes (Lima, Blackwell, DeVault, & Fernández-Juricic, 2015; Neumann et al., 2012). The adverse effects of roads on wildlife, such as habitat fragmentation and habitat loss, are well-known particularly on large carnivores due to their large area requirements, low population densities and direct persecution by humans (Colchero et al., 2011; Crooks, 2002; Grilo, Bissonette, & Santos-Reis, 2009; Stewart et al., 2016). WVCs have been widely recognized as a major impediment to management and

conservation of certain carnivores (Grilo et al., 2009); for instance, the Florida panther (*Puma concolor*; Taylor, Buergelt, Roelke-Parker, Homer, & Rotstein, 2002), the Iberian lynx (*Lynx pardinus*) in Spain (Ferrerias, Gaona, Palomares, & Delibes, 2001), and the ocelot (*Leopardus pardalis*) in Texas (Haines, Tewes, Laack, Grant, & Young, 2005; Hewitt, Cain, Tuovila, Shindle, & Tewes, 1998; Tewes & Hughes, 2001). Furthermore, WVC can act synergistically with other disturbance factors, contributing to increase the fragmentation and isolation of threatened species. For example, Cullen et al. (2016) showed that jaguar (*Panthera onca*) populations are being threatened by a combination of factors, namely habitat loss, habitat fragmentation, and proximity to roads. (Espinoso, Celis, & Branch, 2018).

* Corresponding author.

E-mail address: mkaboli@ut.ac.ir (M. Kaboli).




Fig. 1. Protected areas with cheetah occurrences in Iran.

WVCs are mainly a particular conservation concern for endangered species populations that are already at risk of extinction (Downs et al., 2014). In Iran, WVCs are also well documented for a variety of large mammals including carnivores (Mohammadi & Kaboli, 2016). Among these is one of the most threatened felid species in the world (Farhadinia et al., 2013; Mohammadi & Kaboli, 2016). Prior to the 1940s, it was estimated that around 400 Asiatic cheetahs (*Acinonyx jubatus venaticus*) roamed throughout eastern Iran and in the west near the Iraqi border (Farhadinia et al., 2013). However, during the 2000's, only 82 cheetahs were detected by camera-tracking in their habitats (TBR, Kavir National Park, Khosh Yeilagh Wildlife Refuge, Naybandan Wildlife Refuge, Bafgh Protected Area, Dareh Anjir Wildlife Refuge, Siah Kouh National Park, Ariz No Hunting Area, Abbas Abad Wildlife Refuge, Darband Ravar Wildlife Refuge, Miandasht Wildlife Refuge (Fig. 1)) in Iran (Farhadinia et al., 2017).

Over the past three decades, Iran has been the last refuge for Asiatic cheetahs, occurring throughout several protected areas, including the provinces of North Khorasan, Yazd, Semnan, Kerman and Isfahan (Farhadinia et al., 2017) (Fig. 1). After years of developing road networks in Iran, protected areas with high biodiversity and the corridors that connect them have been severely impacted (Mohammadi & Kaboli, 2016). Yazd and Semnan Provinces in central and northeast of Iran have been among the most problematic areas as a result of expanding anthropogenic activities (including construction of road networks), which leads to human-wildlife conflicts (Mohammadi & Kaboli, 2016).

Road densities within and at the border of the Asiatic cheetah protected areas are shown in the supplementary material (Table S1). Reports indicate that 14 Asiatic cheetahs were killed in vehicle accidents in these regions over the course of 10 years (CACP, 2015) (Fig. 2).

An important step in mitigating the impacts of roads on wildlife is to focus on road-kill hotspots (Litvaitis & Tash, 2008). Road-kill hotspots have typically been identified along roadside habitats attractive to wildlife (e.g., mineral licks, food resources), or wildlife movement corridors (Litvaitis & Tash, 2008). Roads with high traffic volumes and/or high-speed vehicles are associated with increased road-kills (Danks & Porter, 2010; Grilo et al., 2009; Jaarsma & Willems, 2002). Studies suggest that mitigation planning based on sound science can effectively reduce WVCs (Clevenger, Chruszcz, & Gunson, 2001; Rytwinski et al., 2016). As a result, planning and implementation of mitigation measures



Fig. 2. Female Asiatic cheetah killed on the Semnan – Mashhad road due to a vehicle collision in November 2015.

has proliferated worldwide (Beckmann & Hilty, 2010). Unfortunately, few WVC-related studies have been carried out in Asia and even fewer in Iran (Gubbi, Poornesha, & Madhusudan, 2012; Lateef, 2010; Mohammadi & Kaboli, 2016; Moqanaki & Cushman, 2016; Saeki & Macdonald, 2004).

The objectives of our study are the following: 1) collect field data on incidence of WVCs and road crossings in Semnan-Mashhad highway in the northeast of Iran; 2) develop models that identify potential areas of conflict between wildlife movement and road systems; 3) validate models using field data collected; 4) culvert monitoring for wildlife track detection in crossing hotspots; and, 5) recommend mitigation measures to reduce WVCs including mammals, such as Asiatic cheetahs.

2. Material and methods

2.1. Study area

Touran Biosphere Reserve (TBR) is located south of Shahrud and 28 km southeast of Biarjomand, and covers an area of 1,464,992 ha (Fig. 3). In 1976, it was declared as a protected area and in 1977, UNESCO's Man and Biosphere Program (MAB) decided to designate the




Fig. 3. Touran Biosphere Reserve in north east of Iran.

area as a biosphere reserve on account of its unique endemic fauna and flora, which includes a great number of large mammals such as Persian wild ass (*Equus hemionus*), Persian Gazelle (*G. subgutturosa*), Jebeer Gazelle (*G. bennettii*), Jungle cat (*Felis chaus*), Sand cat (*F. margarita*), Leopard (*Panthera pardus*) and most importantly the Asiatic cheetah. TBR is the second largest biosphere reserve in the world and consists of three areas of Wildlife Refuge, National Park, and Protected Area (CACP, 2015). Rangeland is the most dominant ecosystem type in TBR. It is among the most valuable habitats for Asiatic cheetah and appears to be a source for cheetah population in Iran (CACP, 2015). The Miandasht Protected Area, Behkade Protected Area and Golestan National Park are located in the east and north of TBR and act as peripheral core areas for the cheetah population (CACP, 2015). The climate of the region is cold arid (as determined by the Ambrege method). The area encompasses a range of mountains, ridges, plains, sand dunes and sand sheets (CACP, 2015).

The Semnan-Mashhad road is at the border of the TBR, about 45 km of which crosses the northern border of the reserve (Fig. 3). The 12 m wide, 2-lane highway has a traffic volume of 6728 and 719 vehicles per day for private and commercial vehicles, respectively. The speed limit is 90 km/h. The highway has lanes that are separated by concrete and guardrail in some sections of the road (Iran Transportation Organization, 2013). Roadside vegetation is predominantly *Zygo-phyllum eurypterum*, *Artemisia sieberitii* and *Ephedra intermedia* (CACP, 2015).

2.2. Wildlife-vehicle collision (WVC) data collection

We monitored a 20 km section of the Semnan-Mashhad road that borders the TBR. In this 20 km section, we surveyed a 30 m buffer (on either side of the road, corresponding to a 60 m × 20 km area) for road-killed mammals every morning during two periods: late summer (20 August–29 September); and, early autumn (10 October–25 November) of 2014 and 2015. We inspected each side of the road separately on foot and all carcasses of mammals were recorded using a handheld GPS (Garmin GPS Map 62S). In order not to double-count carcasses, we removed the carcasses after recording.

We also obtained WVCs location data from 2011 to 2015 from Semnan province, Department of Environment. We acquired anecdotal data related to mammals crossing the road over a 1-year period in TBR with the contribution of a number of experienced local people who were trained prior to data collection (data included direct observation of mammals in the vicinity of the road and crossing the road). We estimated the probability of a mammal getting killed as it attempts to cross a road, based on an equation used in previous studies (Gibbs & Shriver, 2002; Hels & Buchwald, 2001; Litvaitis & Tash, 2008; Row, Blouin-Demers, & Weatherhead, 2007).

$$P_{killed} = 1 - e^{-Na/v} \quad (1)$$

This equation attempts to ascertain the probability of an animal getting killed (P_{killed}) while crossing the road, given the traffic intensity N (number of vehicles per 24 h) (vehicles/lane/min), the kill zone a (m), and the velocity of the individual v (m/min) while crossing the road (Hels & Buchwald, 2001; Litvaitis & Tash, 2008). In this study, we

considered v : 480 m/min, a : 12 m and n : 4 vehicles/lane/min. We applied the value obtained from this equation to the crossing data

2.3. WVCs hotspots

Here, we determined WVCs hotspots based on two methods, the Distance Method and the Kernel Density Estimation (KDE) technique and compared both methods (Anderson, 2009; Krisp & Durot, 2007; Williamson, McLafferty, Goldsmith, Mollenkopf, & McGuire, 1998; Xie & Yan, 2008).

2.3.1. Distance method

This method was separately applied for each data set (crossing points and WVCs data). As the distance from each point increases, each cell value will gradually increase and the fuzzy raster map will depict the distribution of all points (Quddus, Noland, & Ochieng, 2006; Quinn, Alexander, Heck, & Chernoff, 2011; Wang, Lao, Wu, & Corey, 2010). In this approach, in order to distinguish mortalities of each mammal and to weight each point, we considered two different ecological values for each point: (1) evolutionarily distinct and globally endangered (EDGE); and, (2) home range radius.

2.3.1.1. EDGE. The evolutionarily distinct and globally endangered (EDGE) species is an approach for conservation prioritization of species (Isaac, Turvey, Collen, Waterman, & Baillie, 2007), in which based on the evolutionary distinctiveness and extinction risk according to the equation derived by Isaac et al. (2007), species are prioritized for effective management and conservation (Isaac, Redding, Meredith, & Safi, 2012, Isaac et al., 2007; Jetz et al., 2014).

In this study, we prepared a list of mammals killed on the studied road and prioritized them based on the EDGE method (Isaac et al., 2007). Then, we converted the EDGE values to relative values between 0 and 1. Finally, the relative weight for each mammal was achieved using the relative values (Table 1).

2.3.1.2. Home range radius. This method is based on the home range radius of each mammal, including Persian gazelle (Durmus, 2010), Asiatic cheetah (CACP, 2015), striped hyena (Wagner, Frank, & Creel, 2008), golden jackal (Admasu, Thirgood, Bekele, & Karen Laursen, 2004), red fox (Trehwella, Harris, & McAllister, 1988), European hare (Reitz & Léonard, 1994; Smith, Alves, Ferrand, & Hackländer, 2008), caracal (Avenant & Nel, 1998), and grey wolf (Mattisson et al., 2013). Similar to the previous method, we provided relative values between 0 and 1. Therefore, the relative weight for each mammal can be achieved using the relative values (Table 2).

All procedures of identifying potential hotspot locations followed the following steps: 1) Preparing a raster map using both data sets (crossing points and WVCs) based on Euclidean Distance; 2) Providing a

Table 1
Prioritized list of mammals based on the EDGE method.

	EDGE	IUCN Categories	EDGE/ Total	Relative weight
<i>Acinonyx jubatus venaticus</i>	5.78	CR	0.250	5
<i>Hyaena hyaena</i>	3.76	NT	0.150	3
<i>Capra aegagrus</i>	3.34	VU	0.130	2.6
<i>Ovis orientalis</i>	3.24	VU	0.120	2.4
<i>Gazella subgutturosa</i>	2.72	VU	0.110	2.2
<i>Caracal caracal</i>	2.41	LC	0.100	2
<i>Lepus europaeus</i>	2.01	LC	0.070	1.4
<i>Canis aureus</i>	1.52	LC	0.055	1.1
<i>Vulpes vulpes</i>	1.51	LC	0.055	1.1
<i>Canis lupus</i>	1.45	LC	0.050	1
Total:	21.95		1	

Table 2
Values of each mammal based on home range radius.

	Home range (Km ²)	Radius of home range (Km ²)	Radius of home range / Total	Relative weight
<i>Acinonyx jubatus venaticus</i>	170	7.5	0.280	40
<i>Hyaena hyaena</i>	80	5.2	0.196	28
<i>Canis lupus</i>	70	4.7	0.178	25
<i>Caracal caracal</i>	30	3	0.113	16
<i>Gazella subgutturosa</i>	15	2.1	0.080	11
<i>Ovis orientalis</i>	10	1.7	0.064	9
<i>Vulpes vulpes</i>	3	1	0.038	5.4
<i>Canis aureus</i>	3	1	0.038	5.4
<i>Lepus europaeus</i>	0.7	0.2	0.007	1
Total	26.4	1		

fuzzy map using the raster map to categorize cell values between 0 and 1; 3) Applying EDGE and home range radius values to separately weight mortality points for each map; 4) Inverting each cell value to exhibit both values (EDGE and home range radius) for each point; 5) Overlaying the entire map to prepare hotspot locations and 6) Using contour lines for prioritization of hotspot locations.

2.3.2. Kernel density estimation (KDE)

KDE placed a grid over the WVCs and crossing points; a 10 × 10 m grid cell was used to include the entire distribution of WVC points in the study area. A crucial step in KDE methodology was to determine the proper bandwidth for our study (Anderson, 2009; Krisp & Durot, 2007; Williamson et al., 1998; Xie & Yan, 2008). First, we determined the bandwidth using the nearest neighbor distance algorithm (Williamson et al., 1998), and then computed point weights within the kernel radii. Central point's play a more important role in determining the density value of the cell because the closer a point is to the center, the more weight it will be given. We calculated the final value for each grid cell by summing the values of circle surfaces associated with each point (Silverman, 1986).

2.4. Habitat corridor for Asiatic cheetah

A habitat suitability map of the Asiatic cheetah was prepared using MaxEnt software version 3.3.3k (Phillips, Anderson, & Schapire, 2006) to create habitat patches and apply to corridor modeling. MaxEnt is a general-purpose method for characterizing species habitat associations using presence-only data, and has proven to be robust and precise compared to other methods (Elith et al., 2006). We used 10,000 as the number of pseudo-absence points (Phillips & Dudík, 2008). For the training data set, 75% of the presence points were randomly chosen to produce the model, while the remaining 25% was used as test data (Pearson, Raxworthy, Nakamura, & Peterson, 2007). We used the area under the ROC curve (AUC) to evaluate model performance. AUC ranges from 0.5 to 1, in which 0.5 shows no discrimination between presence points and pseudo-absence points and 1 shows the highest level of discrimination (Phillips et al., 2006).

A total of 365 presence points were obtained from CACP (2015) in TBR and Miandasht Wildlife Refuge, as well as nearby surroundings. Environmental layers, including (1) digital elevation model (DEM) and (2) topographic position map as topographic layers, (3) land cover, (4) distance to river and (5) distance to springs as layers representing food and water needs, (6) distance to road, (7) distance to village and (8) distance to pasture as human disturbance, All layers had a 30 m × 30 m resolution. Topographic position was derived from DEM in four categories (Majka, Jenness, & Beier, 2007); in 200 m radius neighborhood, a 30 × 30 m cell was classified as canyon bottom if cell elevation was at least 12 m lower compared to the neighbourhood average, ridge-top if cell elevation was at least 12 m higher compared to the neighbourhood

average, flat-gentle slope if the cell is neither a canyon bottom nor a ridge-top and slope $< 6^\circ$, and steep slope if the cell is neither a canyon bottom nor a ridge-top and slope $> 6^\circ$. Elevation and slope thresholds were chosen based on methods used by Atwood et al. (2011). Land cover maps (FRWMO (Forest, Range & Watershed Management Organization of Iran), 2010) initially including 45 classes were converted to 11 classes based on the similarities between classes. Other maps were created by the Department of Environment (DoE) of Iran. Distance to these feature maps was calculated using the Euclidean distance tool in the Spatial Analyst extension of ArcGIS 10.2. The habitat suitability map created by MaxEnt was rescaled from a 0–1 range to the 0–100 range.

A habitat suitability map was used to provide potential population patches and potential stepping stone patches (Majka et al., 2007). A 200 m radius neighborhood was chosen to assess each cell of the habitat suitability map. Potential stepping-stone patches were selected based on at least 10 km² of contiguous cells with a neighborhood suitability of > 30 . A threshold of 30 was chosen based on the equal training sensitivity and specificity logistic threshold in MaxEnt results (Liu, Berry, Dawson, & Pearson, 2005) and a threshold of 10 km was chosen based on home range of the Asiatic cheetah (CACP, 2015). Potential population patches were selected based on at least 50 km² of contiguous cells with a neighborhood suitability of > 30 . A 50 km² area is large enough to support about five cheetahs to create a population (CACP, 2015).

Corridor modeling was done using CorridorDesigner software (Majka et al., 2007) in Arc GIS 10.2. The corridor was illustrated between TBR and Miandasht Wildlife Refuge. CorridorDesigner used population patches within each protected area as termini and the corridor was illustrated using the least cost method (Almasieh, Kaboli, & Beier, 2016).

2.5. Culvert usage

Using the track-pad technique, the most appropriate technique in wildlife track detection (Mateus, Grilo, & Santos-Reis, 2011), 10 box-type drainage culverts ($n = 4$, width: $> 2\text{--}3\text{ m} \times$ height: $1\text{--}2.50\text{ m}$; $n = 3$, width: $3\text{--}4.10\text{ m} \times$ height: $2\text{--}3\text{ m}$; $n = 3$, width: $> 4\text{ m} \times$ height $> 3\text{ m}$) in crossing hotspot locations were monitored during winter (November–January) and summer (June–September) of 2015. To detect mammal tracks, we covered the width of all culverts with a thin layer of marble dust, approximately 1 cm deep and 150 cm in length (Mata, Hervas, Herranz, Suarez, & Malo, 2005).

Animal tracks were detected over a 2–3-day interval between visits. Drainage culverts were sampled during seven consecutive days. This sampling period was selected as it is the driest period of the year and when culverts are least likely to be flooded and marbled dust is not washed away.

3. Results

3.1. Wildlife-vehicle collisions

Our results showed that 14 Asiatic cheetah were killed on the roads of Iran during 2005–2016 (CACP, 2015) (Table 3). Among these, seven Asiatic cheetahs were killed on Semnan-Mashhad road. In this study, 71 WVC observations were made of mammalian species, including: Persian gazelle ($n = 5$); Asiatic cheetah ($n = 7$); striped hyena ($n = 4$); golden jackal ($n = 10$); red fox ($n = 20$); European hare ($n = 20$); caracal ($n = 3$); and, grey wolf ($n = 4$). Also, 40 road crossings by mammalian species were recorded, including: Persian gazelle ($n = 9$); grey wolf ($n = 17$); red fox ($n = 9$); and, golden jackal ($n = 5$). Using the nearest neighbor distance algorithm, the bandwidth of 500 m was determined to be the most effective in estimating the density of WVCs. The probability of a mammal getting killed while crossing the road was 0.18 which means 18 percent of the species that cross the road will be killed

Table 3
Road mortalities of the Asiatic cheetah during 2005–2016 (CACP, 2015).

Road	Year	Sex	Season
Calmand - Bahadoran	2005	Female	January
Bafgh- Bahabad	2006	Unknown	December
Calmand - Bahadoran	2006	Male	February
Calmand - Bahadoran	2008	Male	February
Calmand - Bahadoran	2009	Male	October
Calmand - Bahadoran	2009	Female	December
Yazd - Tabas road	2012	Male	May
Semnan - Mashhad road	2011	Female	December
Semnan - Mashhad road	2011	Cub	December
Semnan - Mashhad road	2011	Cub	December
Semnan - Mashhad road	2015	Cub (Female)	November
Semnan - Mashhad road	2015	Female	November
Semnan - Mashhad road	2016	Female	December
Semnan - Mashhad road	2016	Male	May

by vehicles collisions.

Based on the distance method, and considering the EDGE and home range radius values, hotspot locations were determined (Fig. 4), which covered an overall length of 12 km (Fig. 4). Also, when we considered home range radius values of other large mammals such as the grey wolf and the Persian gazelle, vehicle collisions coincided with hotspot locations near Abbas Abad village. The final map of hotspot locations was prioritized based on quantitative values of Euclidean distance cells (Figs. 4 and 5). Asiatic cheetah mortalities are located in hotspot locations (1 and 2) with high priority for urgent monitoring. Another hotspot location (3: medium priority for future monitoring) is located near Abbas Abad village (Fig. 5). Based on KDE, we obtained two hotspot locations (Fig. 6).

3.2. Habitat patches and corridor for Asiatic cheetah

Our MaxEnt model showed an AUC of 0.97 which represents high accuracy of the model. Two modeled population patches were obtained from the habitat suitability model; each patch within each protected area (Fig. 7). The population patches within TBR and Miandasht Wildlife Refuge were about 3447 km² and 996 km², respectively. There were four modeled stepping-stone patches between the two protected areas, covering an area of about 153 km².

The corridor between TBR and Miandasht Wildlife Refuge is shown in Fig. 7. This corridor is about 55 km length and 656 km² area, which connects two population patches within these protected areas. Asiatic cheetah vehicle collisions ($n = 7$) and other mammal collisions such as golden jackal ($n = 7$), red fox ($n = 5$), European hare ($n = 10$) occurred where a road crossed this corridor at the border of the TBR.

3.3. Use of culverts by mammals

Most crossings were associated with the red fox, golden jackal, European hare, the grey wolf and the striped hyena (Table 4). Two records of Asiatic cheetah crossing were reported in medium and large culverts ($n = 1$, width: $2.20\text{ m} \times$ height: $1.60\text{ m} \times$ length: 12 m ; $n = 2$, width: $3.10\text{ m} \times$ height: $2.70\text{ m} \times$ length: 12 m) (Table 4, Fig. 8). The majority of crossings were detected in medium and large box culverts (Table 4).

4. Discussion and conclusions

Populations of the Asiatic cheetah are distributed over a wide area in Iran and individuals constantly travel across vast distances through central parts of the country (in four parks and reserves including Dareh-Anjir Wildlife Refuge, Ariz Non-hunting Area, Bafq Protected Area, and Siah-Kouh National Park). Hence, populations of this subspecies are forced to get across several roads within these protected areas




Fig. 4. Hotspot locations of WVCs based on the distance method with reference to EDGE and home range radius values on Semnan-Mashhad road (high values (red zones) indicate highly significant locations of WVCs hotspots and low values (yellow zones) indicate locations of minor significance in terms of WVCs). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

(Farhadinia et al., 2013).

The population patch within TBR was about 3.5 times larger than the population patch in Miandasht Wildlife Refuge, an indication of better habitat suitability for the Asiatic cheetah in TBR. In other words, contiguous cells with suitability > 30 were 3.5 times higher in TBR than Miandasht Wildlife Refuge. Two of the four stepping-stones were included in the modeled corridor, which covered about 11% of the corridor area (78 km² stepping-stones of 656 km² corridor). These stepping-stones should facilitate movement of Asiatic cheetah individuals within the corridor, acting as stopovers, particularly for long movements of cheetahs within corridors (Almasieh et al., 2016; Moqanaki & Cushman, 2016). Efforts should be made to maintain or restore habitat within the Asiatic cheetah corridor, ensuring functional habitat connectivity between stepping-stone patches and within the corridor. The two stepping-stone patches outside the corridor with an area of 75 km² could potentially be important habitat should efforts be made to create suitable habitat conditions to improve connectivity with the corridor and in the matrix proper. A regional population conservation strategy needs to focus on reducing cheetah mortality on the Abbas Abad Road, through use of existing culverts and construction of new passages, and ensuring travel habitat exists within the corridor.

Previous studies have shown that high rates of mortality may occur within habitat corridors and well-connected areas (Grilo, Ascensao, Santos-Reis, & Bisonette, 2011). Our results revealed that the modeled corridor and vehicle-collisions of the Asiatic cheetah overlapped significantly (Fig. 7). Indeed, Semnan-Mashhad road has bisected this corridor as well as suitable habitats of the species. Mortalities of other mammals, such as the grey wolf, red fox and golden jackal mostly occurred close to Abbas Abad village road due to their scavenging behavior. In areas where roads are barriers to movement, culverts and

crossing structures may serve as corridors to travel across roads (Lister, Brocki, & Ament, 2015). Today, studying the adverse effects of roads on species mobility and the potential for passage structures in alleviating those impacts is a key conservation issue (Lister et al., 2015; Tigas, Van Vuren, & Sauvajot, 2002). Most studies documented that large mammals, especially large carnivores, used crossing structures in a way that ensured habitat connectivity (Gagnon, Dodd, Ogren, & Schweinsburg, 2011; Sawaya, Clevenger, & Kalinowski, 2013). For this reason, determining the precise location of implementing mitigation measures is important (Beckmann & Hilty, 2010; Clevenger & Wierchowski, 2006; Landguth, Hand, Glassy, Cushman, & Sawaya, 2012).

There are many culverts in the identified hotspot. However, only a few of them have appropriate dimensions for the species to cross. Our road-kill data showed that in hotspot locations (1 and 2) of the Asiatic cheetah, there are six drainage culverts; two of which have high openness ratios (Table 4). Also, our study is the first to report on mammal use of below-grade passage structures in Iran and the first documented passage by Asiatic cheetahs.

Most crossings of drainage culverts were recorded from European hare, red fox, golden Jackal, grey wolf. Two records of the Asiatic cheetah crossings were recorded from medium and large culverts (Table 4). The distance between the two culverts is about 300 m. The floor of both culverts used by Asiatic cheetah had a substrate of soft clay soil, and they are used by livestock only in autumn, which together with the fact that visibility of both culverts is suitable (i.e. Asiatic cheetahs can see the end of culverts) might be promoting its use by cheetahs.

In a study in Banff National Park from 1996 to 2000, 22 crossing structures were monitored year-round along 45 km stretch of a highway (Gloyne & Clevenger, 2001). Their results showed that cougars preferred to cross open-span underpasses compared to other types of




Fig. 5. Prioritization of WVCs hotspot locations on Semnan-Mashhad road (Locations 1 and 2 show high-priority hotspots, locations 3 and 4 show medium-priority hotspots for future monitoring).

crossing structures. Moreover, they found that crossing structure use by cougars (including underpasses) was highest during winter and lower in summer, and largely based on seasonal altitudinal migrations and habitat use of their prey species. In our study, the Asiatic cheetah used culverts during both summer and winter seasons, suggesting that their prey and food resources are available year-round. Some large carnivores such as grey wolves prefer large, open structures with good visibility, for instance landscape bridges, wildlife overpasses or viaducts (Clevenger & Waltho, 2005). Red foxes and golden jackals generally experience high levels of road-related mortality (Markolt, Szemethy, Lehoczki, & Heltai, 2012). Studies showed that to encourage red fox and golden jackal use of underpasses and culverts, they should be designed relative to the species' body size (Grilo et al., 2009; Markolt et al., 2012). Other studies showed that crossing structures and the proximity of suitable habitats disposed these species to vehicle collisions (Grilo et al., 2009; Markolt et al., 2012).

Small and medium-sized mammals, particularly prey species, tend to use passages of a size that allow for their movement, but may limit movement of their larger predators (Clevenger et al., 2001). Construction of wildlife crossings along fenced roads can provide safe road conditions for drivers and mitigate barrier effects on entire wildlife communities (Foster & Humphrey, 1995; Rytwinski et al., 2016). However, rarely is a single crossing structure design effective for a wide range of wildlife species, although due to their large size, viaducts and land bridges meet community level needs (Clevenger & Huijser, 2011; Lister et al., 2015). Wildlife species are capable of using a wide range of crossing structure types (Clevenger & Waltho, 2005; Grilo, Bissonette, & Santos-Reis, 2008; Mata et al., 2005). Our data bear this out as well (Table 4). Asiatic cheetahs used small and medium-sized underpasses, red foxes, golden jackals and striped hyenas used medium-sized underpasses, and wolves used mostly large passage structures. Therefore,

a wide range of wildlife crossing types and designs should be used in order to facilitate passage by most species in the affected area. By placing a mix of crossing design types in the 12 km hotspot area, it will help reduce mortality and maintain connectivity for cheetahs and other species we documented in this study area.

Road networks can create suitable habitats for wildlife with a higher availability of food compared to the surrounding environments (Meunier, Verheyden, & Jouventin, 2000; Munguira & Thomas, 1992). Due to our field experiment during winter, when vegetation cover is abundant along the edge of the road, especially in plain areas, Persian gazelles are drawn to these edges, leading to an increase mortality risk from vehicles. Most mortalities of Persian gazelles occurred near Abbas Abad village. In this regard, due to the increase in mortality of this species during critical seasons, mitigation method such as roadside vegetation clearance could be considered as an effective measure in reducing mortality rates. When certain mammals forage along the edge of high-traffic roads, there is an unavoidable risk of vehicle collisions, which provides abundant carrion and attracts scavengers to the road area (Clevenger & Huijser, 2011). Striped hyena, red fox and golden jackal are scavengers and feed on the remains of domestic and other animals that have died due to road accidents (Qarqaz, Abu Baker, & Amr, 2004). Therefore, roads can attract scavengers because of the possibility of the presence of dead animals (Knight & Kawashima, 1993; Santos, Carvalho, & Mira, 2011; Tourani, Moqanaki, & Kiabi, 2012). In view of this behavior, most mortalities associated with these species in this area occurred near Abbas Abad village.

Wildlife warning signs are meant to control WVCs by increasing drivers' awareness and thus reducing vehicle speed (Huijser, Mosler-Berger et al., 2015). Most studies on the effectiveness of signs in reducing WVCs could not ensure their effectiveness (e.g., Bullock, Malan, & Pretorius, 2011; Coulson, 1982; Meyer, 2006; Pojar, Prosenice, Reed,




Fig. 6. Hotspot locations of WVCs based on Kernel Density Estimation with discriminate presence points and WVCs.




Fig. 7. Modeled population patches, modeled stepping stone patches and the modeled corridor between TBR and Miandasht wildlife refuge.

Table 4
Attributes of 10 culverts used by mammals in Touran Biosphere Reserve.

Culvert Attribute	Culverts									
	1	2	3	4	5	6	7	8	9	10
Width	2.20	2.40	2.70	2.80	3.10	3.50	4.10	4.20	4.30	4.30
Height	1.60	1.80	1.90	2.50	2.70	2.90	3.00	3.10	3.35	3.45
Length	12	12	12	12	12	12	11	11	12	12
Openness	0.29	0.36	0.42	0.58	0.69	0.84	1.11	1.18	1.2	1.2
Species										
Red fox	0	0	1	1	1	20	40	0	0	0
Golden Jackal	0	0	0	1	2	12	25	10	0	0
Grey wolf	0	0	0	0	2	3	5	6	8	4
European hare	0	0	1	3	2	5	0	5	0	0
Striped hyena	0	0	0	0	1	1	2	0	0	0
Asiatic cheetah	1	0	0	0	1	0	0	0	0	0
Total	1	0	2	5	9	41	72	21	8	4



Fig. 8. The Asiatic cheetah has crossed the medium culvert (width: 2.20 m, height: 1.60 m, length: 12 m).



Fig. 9. Asiatic cheetah warning sign designed by the CACP project.

& Woodard, 1975; Rogers, 2004). One study found warning signs reducing WVCs by 34% promptly after installment (Found & Boyce, 2011). Other studies have used signage at fence openings where a

crosswalk was painted on the road and resulted in 37–43% decreases (Huijser, Kociolek et al., 2015). Animal-detection systems with or without fencing have proven most effective recently at reducing WVCs by > 50% (Huijser, Kociolek et al., 2015).

The CACP project designed eight Asiatic cheetah warning signs (Fig. 9), for increasing public awareness. Cheetah mortality rates and critical location must be assessed before installment of signs in order to measure their effectiveness in reduction of Asiatic cheetah vehicle collision. Moreover, wildlife fencing and crossing, if designed and maintained properly, are able to minimize vehicle collisions significantly (80–97%) (Huijser et al., 2016). Studies showed that wildlife fencing implemented along short road sections (< 5 km) were less effective in reducing collisions with large mammals than fencing implemented along long road sections (> 5 km) (Huijser et al., 2016). In our study total length of hotspots location was 12 km. In this section there are culverts with suitable dimensions for use by large carnivores, but some culverts are too small, so fencing with plans to retrofit culverts is recommended. Once these structures are retrofitted, systematic monitoring over a minimum of five years should produce useful information to guide the design of crossings for Asiatic cheetahs in future transportation projects in their range. Moreover, use of fencing in hotspot locations might be needed to guide mammals toward the crossing structures. Also, maintenance and monitoring of the crossings should be considered for effective mitigation measures.

In our study, we determined hotspot locations for mammals, particularly

the endangered Asiatic cheetah. We suggest that future studies examine road and environmental factors that may best explain why and where collisions occur. There also are critical gaps in our understanding of road impacts on the population viability of Asiatic cheetahs and other mammals in Iran. Addressing these information needs with focused research should be a high priority as transportation infrastructure expands throughout the country. Spatially explicit individual-based modelling approaches can be an important conservation tool to evaluate how roads affect the population viability of Asiatic cheetahs and other mammals (Grilo et al., 2009; Paula & Desbiez, 2013). This information will help create predictive models of mortality risk and population fragmentation that will identify where Asiatic cheetahs are most threatened by new and improved road systems in Iran. This science-based information will help transportation practitioners in Iran make informed decisions regarding transportation planning that will minimize impacts on Asiatic cheetahs and other sensitive species.

Conflict of interest

The authors declare that they have no conflict of interest.

Acknowledgments

We would like to thank the Conservation of the Asiatic Cheetah Project (CACP), Iranian Department of the Environment (DoE), Yazd and Semnan Provincial Office of DoE particularly Rajabali Karegar and Javad Najafi who provided generous support for field surveys and data collections. We thank two anonymous reviewers for their helpful comments on improving the clarity and content of our manuscript.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.jnc.2018.02.011>.

References

Admasu, E., Thirgood, S. J., Bekele, A., & Karen Laurenson, M. (2004). Spatial ecology of golden jackal in farmland in the Ethiopian Highlands. *African Journal of Ecology*, 42(2), 144–152.

- Almasieh, K., Kaboli, M., & Beier, P. (2016). Identifying habitat cores and corridors for the Iranian black bear in Iran. *Ursus*, 27(1), 18–30.
- Anderson, T. K. (2009). Kernel density estimation and K-means clustering to profile road accident hotspots. *Accident Analysis and Prevention*, 41(3), 359–364.
- Atwood, T. C., Young, J. K., Beckmann, J. P., Breck, S. W., Fike, J., Rhodes, O. E., et al. (2011). Modeling connectivity of black bears in a desert sky island archipelago. *Biological Conservation*, 144, 2851–2862.
- Avenant, N. L., & Nel, J. A. (1998). Home-range use, activity, and density of caracal in relation to prey density. *African Journal of Ecology*, 36(4), 347–359.
- Beckmann, J. P., & Hilty, J. A. (2010). *Connecting wildlife populations in fractured landscapes. Safe passages: highways, wildlife, and habitat connectivity*. Island Press 3–16.
- Bullock, K. L., Malan, G., & Pretorius, M. D. (2011). Mammal and bird road mortalities on the Uppington to Twee Rivieren main road in the southern Kalahari, South Africa. *African Zoology*, 46(1), 60–71.
- CACP (2015). *Conservation of Asiatic Cheetah Project*. Accessed 21-8-16 <http://www.cacp.ir>.
- Clevenger, A. P., & Huijser, M. P. (2011). *Wildlife crossing structure handbook: Design and evaluation in North America (No. FHWA-CFL/TD-11-003)*.
- Clevenger, A. P., & Waltho, N. (2005). Performance indices to identify attributes of highway crossing structures facilitating movement of large mammals. *Biological Conservation*, 121, 453–464.
- Clevenger, A. P., & Wierchowski, J. (2006). *Maintaining and restoring connectivity in landscapes fragmented by roads. Conservation biology series, 14*, Cambridge 502.
- Clevenger, A. P., Chruszcz, B., & Gunson, K. (2001). Drainage culverts as habitat linkages and factors affecting passage by mammals. *Journal of Applied Ecology*, 38, 1340–1349.
- Colchero, F., Conde, D. A., Manterola, C., Chavez, C., Rivera, A., & Ceballos, G. (2011). Jaguars on the move: modeling movement to mitigate fragmentation from road expansion in the Mayan Forest. *Animal Conservation*, 14(2), 158–166.
- Coulson, G. M. (1982). Road-kills of macropods on a section of highway in central Victoria. *Wildlife Research*, 9(1), 21–26.
- Crooks, K. R. (2002). Relative sensitivities of mammalian carnivores to habitat fragmentation. *Conservation Biology*, 16(2), 488–502.
- Cullen, L., Jr., Stanton, J. C., Lima, F., Uezu, A., Perilli, M. L., & Akçakaya, H. R. (2016). Implications of fine-grained habitat fragmentation and road mortality for jaguar conservation in the Atlantic Forest, Brazil. *PLoS One*, 11(12), e0167372.
- Danks, Z. D., & Porter, W. F. (2010). Temporal, spatial, and landscape habitat characteristics of moose - vehicle collisions in Western Maine. *The Journal of Wildlife Management*, 74, 1229–1241. <http://dx.doi.org/10.1111/j.1937-2817.2010.tb01243.x>.
- Downs, J., Horner, M., Loraamm, R., Anderson, J., Kim, H., & Onorato, D. (2014). Strategically locating wildlife crossing structures for Florida panthers using maximal covering approaches. *Transactions in GIS*, 18(1), 46–65.
- Durmus, M. (2010). *Determination of home range size and habitat selection of gazelles (Gazella subgutturosa) by GPS telemetry in Şanlıurfa (doctoral dissertation, Middle East Technical University)*.
- Elith, J., Graham, C. H., Anderson, R. P., Dudík, M., Ferrier, S., Guisan, A., et al. (2006). Novel methods improve prediction of species distributions from occurrence data. *Ecography*, 29, 129–151.
- Espinosa, E., Celis, G., & Branch, L. C. (2018). When roads appear jaguars decline: Increased access to an Amazon wilderness area reduces potential for jaguar conservation. *PLoS One*, 13(1), e0189740.
- Farhadinia, M. S., Akbari, H., Mousavi, S. J., Eslami, M., Azizi, M., Shokouhi, J., ... Hosseini-Zavarei, F. (2013). Exceptionally long movements of the Asiatic cheetah *Acinonyx jubatus venaticus* across multiple arid reserves in central Iran. *Oryx*, 47(03), 427–430.
- Farhadinia, M. S., Hunter, L. T. B., Jourabchian, A., Hosseini-Zavarei, F., Akbari, H., Ziaie, H., et al. (2017). The critically endangered Asiatic cheetah *Acinonyx jubatus venaticus* in Iran: A review of recent distribution, and conservation status. *Biodiversity and Conservation*, 26, 1027–1046.
- Ferreras, P., Gaona, P., Palomares, F., & Delibes, M. (2001). Restore habitat or reduce mortality? Implications from a population viability analysis of the Iberian lynx. *Animal Conservation*, 4(03), 265–274.
- Foster, M. L., & Humphrey, S. R. (1995). Use of highway underpasses by Florida panthers and other wildlife. *Wildlife Society Bulletin (1973–2006)*, 23(1), 95–100.
- Found, R., & Boyce, M. S. (2011). Warning signs mitigate deer-vehicle collisions in an urban area. *Wildlife Society Bulletin*, 35(3), 291–295.
- FRWMO (Forest, Range and Watershed Management Organization of Iran) (2010). *Iranian forests, range and watershed management organization national land use/land cover map*. Accessed 20-6-14 Tehran, Iran: Forest, Range and Watershed Management Organization of Iran. <http://frw.org.ir/00/En/>.
- Gagnon, J. W., Dodd, N. L., Ogren, K. S., & Schweinsburg, R. E. (2011). Factors associated with use of wildlife underpasses and importance of long-term monitoring. *Journal of Wildlife Management*, 75, 1477–1487.
- Gibbs, J. P., & Shriver, W. G. (2002). Estimating the effects of road mortality on turtle populations. *Conservation Biology*, 16(6), 1647–1652.
- Gloyne, C., & Clevenger, A. P. (2001). Cougar *Puma concolor* use of wildlife crossing structures on the Trans-Canada highway in Banff National Park, Alberta. *Wildlife Biology*, 7, 117–124.
- Grilo, C., Ascensao, F., Santos-Reis, M., & Bissonette, J. (2011). Do well-connected landscapes promote road-related mortality? *European Journal of Wildlife Research*, 57, 707–716.
- Grilo, C., Bissonette, J. A., & Santos-Reis, M. (2008). Response of carnivores to existing highway culverts and underpasses: implications for road planning and mitigation. *Biodiversity and Conservation*, 17, 1685–1699.
- Grilo, C., Bissonette, J. A., & Santos-Reis, M. (2009). Spatial-temporal patterns in Mediterranean carnivore road casualties: Consequences for mitigation. *Biological Conservation*, 142(2), 301–313.
- Gubbi, S., Poornesha, H. C., & Madhusudan, M. D. (2012). Impact of vehicular traffic on the use of highway edges by large mammals in a South Indian wildlife reserve. *Current Science*, 102, 1047–1051.
- Haines, A. M., Tewes, M. E., Laack, L. L., Grant, W. E., & Young, J. (2005). Evaluating recovery strategies for an ocelot (*Leopardus pardalis*) population in the United States. *Biological Conservation*, 126(4), 512–522.
- Hels, T., & Buchwald, E. (2001). The effects of road kills on amphibian populations. *Biological Conservation*, 99, 331–340.
- Hewitt, D. G., Cain, A., Tuovila, V., Shindle, D. B., & Tewes, M. E. (1998). *Impacts of an expanded highway on ocelots and bobcats in southern Texas and their preferences for highway crossings. International Conference on Wildlife Ecology and Transportation (ICOWET 1998)*.
- Huijser, M. P., Fairbank, E. R., Camel-Means, W., Graham, J., Watson, V., Basting, P., et al. (2016). Effectiveness of short sections of wildlife fencing and crossing structures along highways in reducing wildlife-vehicle collisions and providing safe crossing opportunities for large mammals. *Biological Conservation*, 197, 61–68.
- Huijser, M. P., Kociolek, A. V., Allen, T. D., McGowen, P., Cramer, P. C., & Venner, M. (2015). *Construction guidelines for wildlife fencing and associated escape and lateral access control measures. Prepared for the American Association of State Highway and Transportation Officials (AASHTO) Standing Committee on the Environment*. p. 218.
- Huijser, M. P., Mosler-Berger, C., Olsson, M., & Strein, M. (2015). Wildlife warning signs and animal detection systems aimed at reducing wildlife-vehicle collisions. In R. Van der Ree, C. Grilo, & D. Smith (Eds.). *Handbook of road ecology* (pp. 198–212). New York, NY: John Wiley & Sons, Ltd.
- Iran Transportation Organization (2013). *National traffic and safety report, (with co-operation with Law Medical Organization)* p. 119.
- Isaac, N. J., Redding, D. W., Meredith, H. M., & Safi, K. (2012). Phylogenetically-informed priorities for amphibian conservation. *PLoS One*, 7(8), e43912.
- Isaac, N. J., Turvey, S. T., Collen, B., Waterman, C., & Baillie, J. E. (2007). Mammals on the EDGE: Conservation priorities based on threat and phylogeny. *PLoS One*, 2(3), e296.
- Jaarsma, C. F., & Willems, G. P. (2002). Reducing habitat fragmentation by minor rural roads through traffic calming. *Landscape and Urban Planning*, 58(2), 125–135.
- Jetz, W., Thomas, G. H., Joy, J. B., Redding, D. W., Hartmann, K., & Mooers, A. O. (2014). Global distribution and conservation of evolutionary distinctness in birds. *Current Biology*, 24(9), 919–930.
- Knight, R. L., & Kawashima, J. Y. (1993). Responses of raven and red-tailed hawk populations to linear right-of-ways. *Journal of Wildlife Management*, 57, 266–271.
- Krisp, J. M., & Durot, S. (2007). Segmentation of lines based on point densities—An optimization of wildlife warning sign placement in southern Finland. *Accident Analysis and Prevention*, 39(1), 38–46.
- Landguth, E., Hand, B., Glassy, J., Cushman, S., & Sawaya, M. (2012). UNICOR: A species connectivity and corridor network simulator. *Ecography*, 35, 9–14.
- Lateef, M. U. (2010). Estimation of fatalities due to road traffic crashes in Karachi, Pakistan, using capture-recapture method. *Asia-Pacific Journal of Public Health*, 22(3), 332–341.
- Lima, S. L., Blackwell, B. F., DeVault, T. L., & Fernández-Juricic, E. (2015). Animal reactions to oncoming vehicles: A conceptual review. *Biological Reviews*. <http://dx.doi.org/10.1111/brv.12093>.
- Lister, N. M., Brocki, M., & Ament, R. (2015). Integrated adaptive design for wildlife movement under climate change. *Frontiers in Ecology and the Environment*, 13, 493–502.
- Litvaitis, J. A., & Tash, J. P. (2008). An approach toward understanding wildlife-vehicle collisions. *Environmental Management*, 42(4), 688–697.
- Liu, C., Berry, P. M., Dawson, T. P., & Pearson, R. G. (2005). Setting thresholds of occurrence in the prediction of species distributions. *Ecography*, 28(3), 385–393.
- Majka, D., Jenness, J., & Beier, P. (2007). *CorridorDesigner: ArcGIS tools for designing and evaluating corridors*. Minneapolis, MN: CorridorDesigner Accessed 25-2-17. <http://www.corridordesign.org/> [Verified 24 January 2017].
- Markolt, F., Szemethy, L., Lehoczki, R., & Heltai, M. (2012). Spatial and temporal evaluation of wildlife-vehicle collisions along the m3 highway in Hungary. *North-Western Journal of Zoology*, 8, 414–425.
- Mata, C., Hervas, I., Herranz, J., Suarez, F., & Malo, J. (2005). Complementary use by vertebrates of crossing structures along a fenced Spanish motorway. *Biological Conservation*, 124, 397–405.
- Mateus, A. R. A., Grilo, C., & Santos-Reis, M. (2011). Surveying drainage culvert use by carnivores: Sampling design and cost-benefit analyses of track-pads vs. video-surveillance methods. *Environmental Monitoring and Assessment*, 181(1–4), 101–109.
- Mattisson, J., Sand, H., Wabakken, P., Gervasi, V., Liberg, O., Linnell, J. D., et al. (2013). Home range size variation in a recovering wolf population: evaluating the effect of environmental, demographic, and social factors. *Oecologia*, 173(3), 813–825.
- Meunier, F. D., Verheyden, C., & Jouvantin, P. (2000). Use of roadsides by diurnal raptors in agricultural landscapes. *Biological Conservation*, 92(3), 291–298.
- Meyer, E. (2006). *Assessing the effectiveness of deer warning signs (No. K-TRAN: KU-03-6)*. Kansas Department of Transportation.
- Mohammadi, A., & Kaboli, M. (2016). Evaluating wildlife-vehicle collision hotspots using kernel-based estimation: A focus on the endangered Asiatic cheetah in central Iran. *Human-Wildlife Interactions*, 10(1), 103–109.
- Moqanaki, E. M., & Cushman, S. A. (2016). All roads lead to Iran: Predicting landscape connectivity of the last stronghold for the critically endangered Asiatic cheetah. *Animal Conservation*, 20(1), 29–41.
- Munguira, M. L., & Thomas, J. A. (1992). Use of road verges by butterfly and burnet populations, and the effect of roads on adult dispersal and mortality. *Journal of Applied Ecology*, 29, 316–329.
- Neumann, W., Ericsson, G., Dettki, H., Bunnefeld, N., Keuler, N. S., Helmers, D. P., et al.

- (2012). Difference in spatiotemporal patterns of wildlife road-crossings and wildlife-vehicle collisions. *Biological Conservation*, 145(1), 70–78.
- Paula, R. C., & Desbiez, A. (2013). Population viability analysis. In A. Consorte-McCrea, & R. Ferraz (Eds.). *Ecology and conservation of the Maned wolf: Multidisciplinary perspectives*. (pp. 15–34). London: CRC Press.
- Pearson, R. G., Raxworthy, C., Nakamura, M., & Peterson, A. T. (2007). Predicting species distributions from small numbers of occurrence records: A test case using cryptic geckos in Madagascar. *Journal of Biogeography*, 34, 102–117.
- Phillips, S. J., & Dudík, M. (2008). Modeling of species distributions with Maxent: New extensions and a comprehensive evaluation. *Ecography*, 31(2), 161–175.
- Phillips, S. J., Anderson, R. P., & Schapire, R. E. (2006). Maximum entropy modeling of species geographic distributions. *Ecological Modelling*, 190, 231–259.
- Pojar, T. M., Prosen, R. A., Reed, D. F., & Woodard, T. N. (1975). Effectiveness of a lighted, animated deer crossing sign. *The Journal of Wildlife Management*, 39(1), 87–91.
- Qarqaz, M. A., Abu Baker, M. A., & Amr, Z. S. (2004). Status and ecology of the striped Hyaena, *Hyaena hyaena*, in Jordan. *Zoology in the Middle East*, 33, 87–92.
- Quddus, M. A., Noland, R. B., & Ochieng, W. Y. (2006). A high accuracy fuzzy logic based map matching algorithm for road transport. *Journal of Intelligent Transportation Systems*, 10(3), 103–115.
- Quinn, M., Alexander, S., Heck, N., & Chernoff, G. (2011). Identification of bird collision hotspots along transmission power lines in Alberta: An expert-based geographic information system (GIS) approach. *Journal of Environmental Informatics*, 18(1), 12–21.
- Reitz, F., & Léonard, Y. (1994). Studies on the European hare. 50. Characteristics of European hare *Lepus europaeus* use of space in a French agricultural region of intensive farming. *Acta Theriologica*, 39(2), 143–157.
- Rogers, E. I. (2004). *An ecological landscape study of deer-vehicle collisions in Kent County, Michigan. Report for the Michigan state police, Office of Highway Safety and Planning*. Amasa, Michigan, USA: White Water Associates, Inc.
- Row, J. R., Blouin-Demers, G., & Weatherhead, P. J. (2007). Demographic effects of road mortality in black ratsnakes (*Elaphe obsoleta*). *Biological Conservation*, 137(1), 117–124.
- Rytwinski, T., Soanes, K., Jaeger, J. A., Fahrig, L., Findlay, C. S., Houlahan, J., et al. (2016). How effective is road mitigation at reducing road-kill? A meta-analysis. *PLoS One*, 11(11), e0166941.
- Saeki, M., & Macdonald, D. W. (2004). The effects of traffic on the raccoon dog (*Nyctereutes procyonoides viverrinus*) and other mammals in Japan. *Biological Conservation*, 118(5), 559–571.
- Santos, S. M., Carvalho, F., & Mira, A. (2011). How long do the dead survive on the road? Carcass persistence probability and implications for road-kill monitoring surveys. *PLoS One*, 6(9), e25383.
- Sawaya, M., Clevenger, A. P., & Kalinowski, S. (2013). Wildlife crossing structures connect Ursid. Populations at wildlife crossing structures in Banff National Park. *Conservation Biology*, 27(4), 721–730.
- Silverman, B. W. (1986). *Density estimation for statistics and data analysis*. London: Chapman & Hall.
- Smith, A. T., Alves, P. C., Ferrand, N., & Hackländer, K. (2008). *Lagomorph biology: Evolution, ecology, and conservation*. Berlin Heidelberg: Springer-Verlag.
- Stewart, F. E., Heim, N. A., Clevenger, A. P., Paczkowski, J., Volpe, J. P., & Fisher, J. T. (2016). Wolverine behavior varies spatially with anthropogenic footprint: Implications for conservation and inferences about declines. *Ecology and Evolution*, 6(5), 1493–1503.
- Taylor, S. K., Buergelt, C. D., Roelke-Parker, M. E., Homer, B. L., & Rotstein, D. S. (2002). Causes of mortality of free-ranging Florida panthers. *Journal of Wildlife Diseases*, 38(1), 107–114.
- Tewes, M. E., & Hughes, R. W. (2001). Ocelot management and conservation along transportation corridors in southern Texas. In G. L. Evinck (Ed.). *Proceedings of the International Conference on Ecology and Transportation* Keystone, Colorado p. 559.
- Tigas, L. A., Van Vuren, D. H., & Sauvajot, R. M. (2002). Behavioral responses of bobcats and coyotes to habitat fragmentation and corridors in an urban environment. *Biological Conservation*, 108(3), 299–306.
- Tourani, M., Moqanaki, E. M., & Kiabi, B. H. (2012). Vulnerability of striped Hyaenas, *Hyaena Hyaena*, in a human-dominated landscape of Central Iran. *Zoology in the Middle East*, 56(1), 133–136.
- Trewhella, W. J., Harris, S., & McAllister, F. E. (1988). Dispersal distance, home-range size and population density in the red fox (*Vulpes Vulpes*): A quantitative analysis. *Journal of Applied Ecology*, 25, 423–434.
- Wagner, A. P., Frank, L. G., & Creel, S. (2008). Spatial grouping in behaviourally solitary striped hyaenas, *Hyaena Hyaena*. *Animal Behaviour*, 75(3), 1131–1142.
- Wang, Y., Lao, Y., Wu, Y.-J., & Corey, J. (2010). *Identifying high-risk locations of animal-vehicle collisions on Washington state highways*. Seattle, USA: Washington State Transportation Center.
- Williamson, D., McLafferty, S., Goldsmith, V., Mollenkopf, J., & McGuire, P. (1998). *Smoothing crime incident data: New methods for determining the bandwidth in Kernel estimation*. 18th ESRI International User Conference, San Diego, 27–31 July <http://gis.esri.com/library/userconf/proc98/PROCEED>.
- Xie, Z., & Yan, J. (2008). Kernel density estimation of traffic accidents in a network space. *Computers, Environment and Urban Systems*, 32(5), 396–406.